

May 20, 2008

K^+ -nucleon interaction

W. R. Gibbs and R. Arceo

Department of Physics,

New Mexico State University

Las Cruces, New Mexico 88003, USA

Abstract

The Born approximation to the quark-gluon-exchange mechanism for K^+N scattering is used as a starting point to generate a potential for this system. The valence quark wave function of the nucleon is generalized from a single Gaussian to a sum of Gaussians in order to have a more flexible representation than previous work. We obtain a potential derived from a valence density given by lattice calculations. By comparing with a recent amplitude analysis it is found that the strength of the quark-gluon based potential needs to be increased by a factor of order 2-4 relative to the normalization given by more traditional values of the governing parameters. The method is used to estimate the change in effective K^+N amplitudes which would result from changes in the valence quark distributions or strength of the interaction which might arise from nuclear medium effects in K^+ scattering from nuclei.

PACS numbers: 13.75.Jz, 25.80.Nv, 12.39.Pn

I. INTRODUCTION

The K^+ meson is useful in the study of the strong interaction because of its simple structure, i.e. it has no light valence antiquark. This feature means that no 3-quark intermediate state can exist, thus removing the most common resonances from the interaction. Although the interaction is weaker than other typical strong interactions it is still substantial and must be treated as such.

One of the principal motivations of this work is to obtain the relationship between the distribution of the constituents of the nucleon and the K^+ -N scattering amplitudes. In Ref. [1] it was suggested that the increase in size of the nucleon in the nucleus could be observed as an alteration in the scattering of the kaons from nuclei. In that case the model used to connect the change in size with scattering was a strongly repulsive well whose radius was allowed to vary. While this simple model may be expected to encompass much of the basic physics, clearly a model based on fundamental degrees of freedom is to be preferred.

Barnes and Swanson [2] treated the K^+ -nucleon interaction due to the exchange of one gluon and two quarks, one of the lowest order process dealing directly with the underlying degrees of freedom. (Barnes, Black and Swanson[3] carried out a similar program for meson-meson scattering but we do not address that sector here). Their calculation of a lowest order exchange must be iterated to obtain a full evaluation of the scattering amplitude. A common way to do this is to treat their result as the Born approximation corresponding to a potential and use a wave equation to generate an approximation to the full amplitude.

Such a program has been very successful in the case of one pion exchange. The one-pion-exchange potential is able to predict many properties of the deuteron [4, 5, 6]. In the pion case the exchange of a single particle gives rise to a local potential. In the present (quark-gluon exchange) case the potential has one term which is local and three others which are non-local.

In the case of one pion exchange, a potential is essential in order to be able to produce the deuteron bound state. In the present work the potential is needed to calculate realistic phase shifts for comparison with data. This potential will then enable us to relate possible changes in valence quark distribution or strengths to the changes in phase shifts which would

result.

Because they included no mechanism that would lead to spin dependence, the results of Ref. [2] were necessarily spin-independent. The experimentally determined partial waves with ℓ greater than zero show a strong spin dependence[7] so that a comparison with these partial waves is not possible and they concentrated their work on the $\ell = 0$ partial wave. We will do the same.

In a later, similar, program N. Black[8] calculated the higher partial waves with quark-gluon exchange including the spin dependence. For $T=1$ he found qualitative agreement except for the $P_{3/2}$ partial wave where the sign was wrong. For $T = 0$ the phases shifts were considerably smaller than the experimental ones, although there was a tendency for an alternation of sign similar to what is observed. In any case, the $\ell = 0$ partial wave dominates the isospin one amplitude at low to moderate energies.

The Barnes and Swanson[2] calculation assumes a Gaussian wave function for the distribution of the quarks in the nucleon. This is a rather restrictive assumption on the shape and does not represent the functional form found in lattice calculations[9, 10]. To repeat their calculation directly with more realistic wave functions would be difficult because of the 9-dimensional integrals involved. With Gaussian wave functions these integrals can be done analytically so that an alternative method of treating this problem is by representing the desired wave function as a sum of Gaussian functions. In this case the integrals can be done term by term but involve cross terms in which the Gaussian parameters in the initial and final states are different. Such an approach requires a derivation of the expressions of Ref. [2] for this more general condition. We carry out this step and then calculate a potential for a wave function consisting of the sum of two Gaussian functions. One could include more terms but at this stage the knowledge of the desired wave function is probably not sufficiently accurate to warrant the effort.

In the calculation of Barnes and Swanson[2] the Born approximation to the amplitude for K^+N scattering is given by

$$A(\mathbf{k}, \mathbf{k}') = \frac{2\gamma m_R}{3} \sum_{i=1}^4 \eta_i w_i F_i(\mathbf{k}, \mathbf{k}') \quad (1)$$

where

$$F_i(\mathbf{k}, \mathbf{k}') = e^{-a_i k^2 + b_i \mathbf{k} \cdot \mathbf{k}'}, \quad m_R = \frac{E_K E_N}{E_K + E_N}, \quad (2)$$

$\mathbf{k}$ and $\mathbf{k}'$ are center of mass momenta, $k = |\mathbf{k}| = |\mathbf{k}'|$, $\gamma \equiv \alpha_s/m_q^2$, E_N is the total energy of the nucleon and E_K is the total energy of the kaon in the center of mass, α_s is the running strong coupling constant and m_q is the light (constituent) quark mass. Expressions for the a_i , b_i , and η_i are given in Ref. [2] for simple harmonic oscillator wave functions. They are functions of ρ , the ratio of light to strange quark (taken to be 0.6), α , the parameter governing the size of the nucleon and β , the parameter determining the size of the kaon through the ratio $g = \alpha^2/\beta^2$.

They considered four diagrams for quark and gluon exchange. For the first diagram the momentum transfer is given by the two exchanged quarks with gluon exchange between them so the resulting interaction is local as indicated by the fact that $a_1 = b_1$. For the other three diagrams this is not the case and the interaction is non-local. Barnes and Swanson calculated an equivalent local potential valid at zero energy only.

The isospin dependence is contained in the weights, w_i , which are given by

$$\{w\} = \left\{0, \frac{1}{6}, 0, \frac{1}{6}\right\} : T = 0 \text{ and } \{w\} = \left\{\frac{1}{3}, \frac{1}{18}, \frac{1}{3}, \frac{1}{18}\right\} : T = 1. \quad (3)$$

One sees that there is a bias toward the isospin unity amplitude being larger than isospin zero. If the contributions from each of the four diagrams were equal (they need not be, of course, and are not) then the $T = 1$ amplitude would be 7/3 of the $T = 0$ amplitude.

It is interesting to examine the low-energy limit of this amplitude. In this case the functions, $F_i(\mathbf{k}, \mathbf{k}')$, become unity. The functions, η_i depend only on g so, in this limit, the absolute size of the kaon and nucleon systems does not enter but their *relative* size remains very important.

In Section II we derive expressions for the potential for arbitrary values of the parameters in each term, in Section III we study the results with the use of various single Gaussian wave functions for the valence quarks, and in Section IV we develop the two-Gaussian form for the wave function. In Section V we give a summary of the formulas needed to calculate the parameters for the two-(or multi-)Gaussian potential, in Section VI we show the basic fit with the two-Gaussian form and study the consequences of a variation of the radius or

strength (as for the case of immersion in nuclear matter). Section VII gives the results of a calculation of the off-shell amplitude which results from the two-Gaussian form for the potential and Section VIII gives some conclusions.

II. DEFINITION OF THE POTENTIAL

The potential will be given in terms of the amplitude by

$$V(\mathbf{r}, \mathbf{r}') = \frac{2\pi}{m_R} \frac{2\gamma m_R}{3} \sum_{i=1}^4 \eta_i w_i F_i(\mathbf{r}, \mathbf{r}') \quad (4)$$

where

$$F_i(\mathbf{r}, \mathbf{r}') = \frac{1}{(2\pi)^6} \int d\mathbf{k} d\mathbf{k}' e^{i\mathbf{k}\cdot\mathbf{r} - i\mathbf{k}'\cdot\mathbf{r}'} F_i(\mathbf{k}, \mathbf{k}'). \quad (5)$$

To get a fully off-shell amplitude which reduces to this form on shell but is a function only of momentum transfer if $a = b$, we write (dropping for the moment the index i)

$$F(\mathbf{k}, \mathbf{k}') = e^{-\frac{a}{2}(k^2 + k'^2) + \mathbf{k}\cdot\mathbf{k}'b} \quad (6)$$

where $\mathbf{k}$ and $\mathbf{k}'$ are independent and are assumed not to be constrained by the on-shell condition. We hasten to point out that this is a choice and that other dependences are possible. We make no claim to uniqueness. We now choose new vector variables as

$$\boldsymbol{\kappa} \equiv \frac{\mathbf{k} + \mathbf{k}'}{2}; \quad \mathbf{q} \equiv \mathbf{k} - \mathbf{k}' \quad \text{so that} \quad \mathbf{k} = \boldsymbol{\kappa} + \frac{\mathbf{q}}{2}; \quad \mathbf{k}' = \boldsymbol{\kappa} - \frac{\mathbf{q}}{2} \quad (7)$$

allowing us to express the form as

$$F(\boldsymbol{\kappa}, \mathbf{q}) = e^{-\kappa^2(a-b)} e^{-q^2(a+b)/4}. \quad (8)$$

For the special case of $a = b$ we can take the Fourier transform on $\mathbf{q}$ to get the local potential

$$F_1(r) = \int d\mathbf{q} e^{i\mathbf{q}\cdot\mathbf{r}} e^{-\frac{1}{2}a_1 q^2} = \left(\frac{2\pi}{a_1}\right)^{\frac{3}{2}} e^{-\frac{1}{2}r^2/a_1}. \quad (9)$$

In the general case we can write the Fourier transform as

$$F(\mathbf{r}, \mathbf{r}') = \frac{1}{(2\pi)^6} \int d\mathbf{k} d\mathbf{k}' e^{-i\mathbf{k}\cdot\mathbf{r}} e^{i\mathbf{k}'\cdot\mathbf{r}'} e^{-\frac{a}{2}(k^2 + k'^2) + \mathbf{k}\cdot\mathbf{k}'b} \quad (10)$$

$$= \frac{1}{(2\pi)^6} \int d\mathbf{\kappa} d\mathbf{q} e^{-i\mathbf{\kappa} \cdot (\mathbf{r} - \mathbf{r}')} e^{-i\mathbf{q} \cdot (\mathbf{r} + \mathbf{r}')/2} e^{-\kappa^2(a-b)} e^{-q^2(a+b)/4}. \quad (11)$$

We use the formula

$$\int d\mathbf{r} e^{i\mathbf{q} \cdot \mathbf{r}} e^{-\alpha^2 r^2} = \frac{\pi^{\frac{3}{2}}}{\alpha^3} e^{-\frac{q^2}{4\alpha^2}} \quad (12)$$

to do the two Gaussian integrals separately to obtain

$$F(\mathbf{r}, \mathbf{r}') = \frac{1}{(2\pi)^3 \sqrt{a^2 - b^2}^3} e^{-\frac{(\mathbf{r} - \mathbf{r}')^2}{4(a-b)}} e^{-\frac{(\mathbf{r} + \mathbf{r}')^2}{4(a+b)}} \quad (13)$$

$$= \frac{4\pi}{(2\pi)^3 \sqrt{a^2 - b^2}^3} e^{-\frac{a(r^2 + r'^2)}{2(a^2 - b^2)}} \sum_{\ell=0}^{\infty} Y_{\ell}^{m*}(\hat{\mathbf{r}}) Y_{\ell}^m(\hat{\mathbf{r}}') Q_{\ell} \left(\frac{br r'}{a^2 - b^2} \right) \quad (14)$$

where

$$Q_{\ell}(z) = (-i)^{\ell} j_{\ell}(iz) \quad (15)$$

and $j_{\ell}(iz)$ is the spherical Bessel function of imaginary argument. The local potential can be recovered by taking the limit $b \rightarrow a$ in the non-local expression.

The wave equation to be solved is

$$\nabla^2 \psi(\mathbf{r}) + \frac{2m_R}{\hbar^2} V_1(r) \psi(\mathbf{r}) + \frac{2m_R}{\hbar^2} \sum_{i=2,3,4} \int d\mathbf{r}' V_i(\mathbf{r}, \mathbf{r}') \psi(\mathbf{r}') = k_0^2 \psi(\mathbf{r}) \quad (16)$$

from which the solution for the wave function and phase shifts can be obtained by standard matrix techniques.

Because local potentials are relative easy to deal with, a commonly used approximation has been to create a local potential which is energy dependent although it is inconvenient due to the loss of orthogonality of wave functions at different energies. We considered such a procedure in this case. Thus we could write

$$F(\mathbf{k}, \mathbf{k}') = e^{-\frac{a-b}{2}(k^2 + k'^2) - \frac{b}{2}(\mathbf{k} - \mathbf{k}')^2} \longrightarrow e^{-(a-b)k_0^2 - \frac{b}{2}(\mathbf{k} - \mathbf{k}')^2} \quad (17)$$

where k_0 is the free-space wave number. However, for the standard parameters given in Ref. [2] for the fourth diagram, the value of b_4 is negative so one cannot create a local potential for this term in this way. We then abandoned this procedure and continued with the non-local potential for the terms beyond the first. Of course, an energy-dependent local potential can be generated for the sum of the four terms after the local/non-local system has been solved.

Set	Description	α (GeV/c)	$R(N)$ (fm)	β (GeV/c)	$R(K)$ (fm)	F	Born a_0, a_1 (fm)	Potential a_0, a_1 (fm)
a	Ref. set	0.40	0.85	0.35	0.49	1.00	-0.12, -0.35	-0.10, -0.23
b	Fit	0.68	0.50	0.43	0.40	0.98	-0.15, -0.31	-0.11, -0.18
a'	Mod. Ref.	0.40	0.85	0.50	0.34	2.06	-0.16, -0.78	-0.12, -0.34
b'	Mod. Fit	0.68	0.50	1.15	0.15	4.84	-0.21, -1.89	-0.11, -0.31
c	Lattice	0.61	0.56	0.93	0.18	3.36	-0.18, -1.32	-0.11, -0.31

TABLE I: Parameter values for the single-Gaussian fit to the quark density. $R(N)$ is the rms value of the inter-quark distance in the nucleon and $R(K)$ is the same quantity for the kaon. Also given is the original reference set as well as the fitted set of Ref. [2]. Sets a' and b' are the result of holding the nucleon size fixed at the unprimed values and adjusting β and F to fit the phase shifts.

III. RESULTS FOR SINGLE GAUSSIAN

Many different assumptions have been made about the spacial extent of the valence quark distribution of the nucleon so that we need to consider a possible range of models. One approach which should be useful is the calculation on the lattice. We will use that of Lissia *et al.*[9] although the results of Alexandrou *et al.*[10] are similar. Figure 1 shows (dotted line) the result of a fit of single Gaussian to the data from Lissia *et al.*[9]. For this single-Gaussian case we can use the equations from Ref. [2] directly to compute the amplitude. Since the light quark mass and the strong coupling constant enter only in the combination, $\gamma = \alpha_s/m_s^2$, rather than consider changing these quantities independently we calculate relative to a standard value of $\gamma_0 = 0.6/0.33^2 = 5.51 \text{ (GeV/c}^2\text{)}^{-2}$. We define $F = \gamma/\gamma_0$ and keep the ratio of light to strange quark fixed at $\rho = 0.6$.

Reference [2] first calculated with a “reference” set of parameters taken from quark systematics. For this “reference” set we find that, although the scattering lengths for the Born approximation are in agreement with the amplitude analysis[7], the potential solution is not. The phase shifts from this calculation are shown in Fig. 2. Although the Born result has the correct ratio of iso-triplet to iso-singlet scattering length, the potential result does not. Since




FIG. 1: One and two Gaussian fit to the proton wave function. The points are from Lissia *et al.*[9].

this ratio is sensitive to the relative hadron sizes we can change the kaon size parameter to $\beta = 0.5$ GeV/c to get the correct ratio in the potential calculation and increase the strength of the potential by a factor of 2.06 to get the right magnitude. In this manner a good fit for the potential calculation is found (see case a' in Table I).

If we use the “fitted” values obtained by Ref. [2] we find case b in Table I. For this value of the nucleon size we need to alter the strength of the potential by a factor of 4.84 and choose $\beta = 1.15$ GeV/c to get the correct values for the scattering lengths (Table I).

If we fix the value of $\alpha = 0.61$ GeV/c from the single Gaussian fit to Lissia *et al.*, increase the strength of the potential by a factor of 3.36 and take $\beta = 0.93$ GeV/c we are able to fit the scattering phase shifts well (case c in Table I). Figure 3 shows the result of this calculation along with the “reference” set of Ref. [2].

Figure 4 show the results for the individual diagram potentials without isospin factors from Eq. 3 (equivalent local potential at zero energy for all but the first) for the “reference”,




FIG. 2: Results of Barnes and Swanson with their reference and fitted set in Born approximation (dashed lines) and the corresponding phase shifts from the solution with the potential (solid lines). The phase shifts from Ref. [7] are shown by the solid dots.

“fit” and lattice input of case c . Also shown (short dash curve) is the result of changing α from 0.61 GeV/c to 0.55 GeV/c ($R_{rms} = 0.56$ fm to $R_{rms} = 0.62$ fm).

It is seen that the ranges of the potentials from the second and fourth diagrams (the only ones which contribute to the $T = 0$ amplitude) appear to decrease with increasing nucleon radius whereas the range of the potential from the first and third (which dominate the isospin one amplitude) increase. One can trace this behavior back to the quantities η_i , a_i , and b_i . In fact, the “true” range parameters (a_i , b_i) always increase with increasing R_{rms} but η_2 and η_4 decrease rapidly.




FIG. 3: S-wave phase shifts using a wave function with a single value of α obtained from the fit to Lissia et al. [9]. Born approximation (short dashed lines) and the corresponding phase shifts from the solution with the potential (solid lines). The result of the reference calculation and its potential corrected values are shown by the long dash and dash dot curves respectively. The phase shifts from Ref. [7] are shown by the solid dots.

Figure 5 shows the full potentials for the two isospins for case *c*. It is seen that the potential is very near to a hard-core potential for case *c*. This result can no doubt be traced to the fact that for both isospins the phase shifts are nearly linear as a function of the center-of-mass momentum as is the case for a hard-sphere interaction. Also shown (short dash curve) is the result for $\alpha = 0.55$ GeV/c. As expected from the remarks above, the isospin zero potential has the opposite variation with nucleon size as the isospin one




FIG. 4: Potentials without the isospin factors for K^+N scattering for three of the parameter sets given in Table I. The upper left hand panel shows the local potential corresponding to the first diagram and the rest show the corresponding equivalent local potentials at zero energy. The short-dash curve with c shows the effect of changing α from 0.61 to 0.55 GeV/c.

potential.

These results show that the Born approximation (especially for $T = 1$) is not accurate. The use of a potential (or other unitarizing agent) is essential. The range of the potential (related to the size of the hadronic systems) is important. For a more extended potential the Born approximation is more nearly correct.




FIG. 5: The full isospin potentials for a single Gaussian at zero energy using the equivalent local forms for the non-local potentials calculated with the parameter set c in Table I (solid curve) and with $\alpha = 0.55$ GeV/c (dashed curve).

IV. TWO GAUSSIAN FORM OF THE WAVE FUNCTION

With the extension of the formulas of Ref. [2] given in the next section one could carry out the calculation of a potential using a sum of any number of Gaussian functions for the quark wave function of the nucleon. A similar program was carried out for meson-meson scattering by Hilbert *et al.*[11]. Of course, such a sum would introduce a large number of parameters which have to be determined. Since we intend to fit them to lattice data, and those calculations are of a limited accuracy at present, we use a sum of only two such functions. The wave functions below are expressed in Jacobian coordinates as discussed in Appendix A.

We take the sum of two Gaussian wave functions, each normalized independently. With

$$\psi(p, q) = N_a \phi_a(p, q) + N_b \phi_b(p, q) \quad (18)$$

we have for the normalization integral

$$\int d\mathbf{p} d\mathbf{q} \psi^2(p, q) = 27(N_a^2 + N_b^2) + 2N_a N_b \int d\mathbf{p} d\mathbf{q} \phi_a(p, q) \phi_b(p, q) = 1. \quad (19)$$

Since

$$\int d\mathbf{p} d\mathbf{q} \phi_a(p, q) \phi_b(p, q) = \frac{3^{\frac{3}{2}} (4\pi)^2}{\pi^3 \alpha_a^3 \alpha_b^3} \int p^2 dp q^2 dq e^{-\frac{p^2}{4} \left(\frac{1}{\alpha_a^2} + \frac{1}{\alpha_b^2} \right)} e^{-\frac{q^2}{3} \left(\frac{1}{\alpha_a^2} + \frac{1}{\alpha_b^2} \right)} = 27 \frac{8\alpha_a^3 \alpha_b^3}{(\alpha_a^2 + \alpha_b^2)^3}, \quad (20)$$

the normalization condition becomes

$$N_a^2 + \frac{16N_a N_b \alpha_a^3 \alpha_b^3}{(\alpha_a^2 + \alpha_b^2)^3} + N_b^2 = 1. \quad (21)$$

If we take $N_b = X N_a$ then

$$N_a^2 \left[1 + \frac{16X \alpha_a^3 \alpha_b^3}{(\alpha_a^2 + \alpha_b^2)^3} + X^2 \right] = 1. \quad (22)$$

We are free to take X , α_a and α_b as we wish and, as long as this condition is satisfied, the wave function will be normalized.

V. SUMMARY OF FORMULAS

In order to carry out the calculation of the potential (or the amplitude) for a sum of Gaussian forms for the wave function we must evaluate the integrals given in Ref. [2] (Eqs. 41, 42, 43 and 44 which are our equations B5, B16, B28 and B39) for the case where kaons A and C have different governing Gaussian parameters (β) and nucleons B and D also have different parameters (α). The evaluation of the integrals is done in appendix B and here we give a summary of the results. Auxiliary definitions are given at the end.

$$a_1 = \frac{1}{2\tilde{\beta}^2(1+\rho)^2} + \frac{1}{3\tilde{\alpha}^2}; \quad b_1 = a_1; \quad \eta_1 = \frac{\alpha_1^3 \alpha_2^3 \beta_1^{\frac{3}{2}} \beta_2^{\frac{3}{2}}}{\tilde{\alpha}^6 \tilde{\beta}^3} \quad (23)$$

$$a_2 = \frac{1}{2(1+\rho)^2 \tilde{\beta}^2} + \frac{1}{12\tilde{\alpha}^2} + \frac{C_2^2 + C_2'^2}{4(\alpha_1^2 + 2\beta_{12}^2 + \frac{1}{3}\alpha_{12}^2)} \quad (24)$$

$$b_2 = \frac{1}{2(1+\rho)^2 \tilde{\beta}^2} + \frac{1}{12\tilde{\alpha}^2} - \frac{C_2 C_2'}{2(\alpha_1^2 + 2\beta_{12}^2 + \frac{1}{3}\alpha_{12}^2)} \quad (25)$$

$$C_2 = -\frac{[(2+\gamma_1^\alpha)(1+\rho) - 6\gamma_2^\beta]}{3(1+\rho)}; \quad C_2' = \frac{[-\gamma_1^\alpha(1+\rho) + 6(\gamma_2^\beta + \rho)]}{3(1+\rho)}$$

$$\eta_2 = \frac{2^{\frac{3}{2}} \alpha_1^3 \alpha_2^3 \beta_1^{\frac{3}{2}} \beta_2^{\frac{3}{2}}}{\tilde{\beta}^3 \tilde{\alpha}^3 (\alpha_1^2 + 2\beta_{12}^2 + \frac{1}{3}\alpha_{12}^2)^{\frac{3}{2}}} \quad (26)$$

$$a_3 = \frac{1}{3\tilde{\alpha}^2} + \frac{3(C_3^2 + C_3'^2)}{8(2\alpha_{12}^2 + 3\beta_1^2)}; \quad b_3 = \frac{1}{3\tilde{\alpha}^2} - \frac{3C_3 C_3'}{4(2\alpha_{12}^2 + 3\beta_1^2)} \quad (27)$$

$$C_3 = \frac{2[2\gamma_2^\alpha + (2\gamma_2^\alpha - 3)\rho]}{3(1+\rho)}; \quad C_3' = -\frac{2}{3}(2\gamma_2^\alpha + 1) \quad (28)$$

$$\eta_3 = \rho \frac{6^{\frac{3}{2}} \alpha_1^3 \alpha_2^3 \beta_1^{\frac{3}{2}} \beta_2^{\frac{3}{2}}}{\tilde{\alpha}^6 (2\alpha_{12}^2 + 3\beta_1^2)^{\frac{3}{2}}} \quad (29)$$

$$a_4 = \frac{(1-\rho)^2}{4(\tilde{\alpha} + 2\tilde{\beta})(1+\rho)^2} + \frac{1}{12\tilde{\alpha}^2} + \frac{C_4^2 + C_4'^2}{4\left(\frac{v^2 w^2}{v^2 + w^2} + \frac{1}{3}\alpha_{12}^2\right)} \quad (30)$$

$$b_4 = \frac{(1-\rho)^2}{4(\tilde{\alpha} + 2\tilde{\beta})(1+\rho)^2} + \frac{1}{12\tilde{\alpha}^2} - \frac{C_4 C_4'}{2\left(\frac{v^2 w^2}{v^2 + w^2} + \frac{1}{3}\alpha_{12}^2\right)} \quad (31)$$

$$C_4 = -\frac{2w^2 - v^2 + \frac{6a^2 \rho}{1+\rho} + (v^2 + w^2)\gamma_1^\alpha}{3(v^2 + w^2)}; \quad C_4' = -\frac{\frac{6w^2 \rho}{1+\rho} + 3v^2 - (v^2 + w^2)\gamma_1^\alpha}{3(v^2 + w^2)} \quad (32)$$

$$\eta_4 = \rho \alpha_1^3 \alpha_2^3 \beta_1^{\frac{3}{2}} \beta_2^{\frac{3}{2}} \left(\frac{8}{(v^2 + w^2)\tilde{\alpha}^2} \right)^{\frac{3}{2}} \frac{1}{\left(\frac{v^2 w^2}{v^2 + w^2} + \frac{1}{3}\alpha_{12}^2\right)^{\frac{3}{2}}} \quad (33)$$

$$v^2 = \alpha_1^2 + 2\beta_2^2; \quad w^2 = \alpha_2^2 + 2\beta_1^2 \quad (34)$$

$$\gamma_i^\beta = \frac{\beta_i^2}{\beta_1^2 + \beta_2^2}; \quad \gamma_i^\alpha = \frac{\alpha_i^2}{\alpha_1^2 + \alpha_2^2}; \quad (35)$$

$$\tilde{\alpha}^2 = \frac{1}{2}(\alpha_1^2 + \alpha_2^2); \quad \tilde{\beta}^2 = \frac{1}{2}(\beta_1^2 + \beta_2^2); \quad \alpha_{12}^2 = \frac{\alpha_1^2 \alpha_2^2}{\alpha_1^2 + \alpha_2^2} \quad \beta_{12}^2 = \frac{\beta_1^2 \beta_2^2}{\beta_1^2 + \beta_2^2} \quad (36)$$




FIG. 6: S-wave phase shifts with α_a , α_b and x chosen from Lissia et al. [9] and the overall factor on the potential chosen to fit the $T=1$ phase shift. The dashed curve shows the result if the value of α_a is altered from its fitted value corresponding to the changes in density shown in Fig. 7. The phase shifts from Ref. [7] are shown by the solid dots.

VI. RESULTS FOR A DOUBLE GAUSSIAN WAVE FUNCTION

Figure 1 shows a fit (solid line) of the two-Gaussian type to the lattice calculation of Lissia *et al.*[9]. The values of the double Gaussian parameters are $\alpha_a = 0.58$ GeV/c, $\alpha_b = 1.32$ GeV/c and $X = 0.1$. This selection of parameters gives a reasonable representation of the s-wave phase shift (see Fig. 6) but a factor of 4.1 has been multiplied by the potential to obtain the fit.




FIG. 7: The solid curve gives the result for $r^2\rho(r)$ with α_a , α_b and X chosen from Lissia et al. [9]. The curves show the densities that result for various values of α_a with $\alpha_b = 1.32$ GeV/c and X fixed at 0.1.

The fit with the double Gaussian is shown in Fig. 6 as the solid line. Also shown are a series of variations of α_1 (holding α_2 and X fixed) used to vary the radius of the system. The effect of this variation on the density is shown in Fig. 7. This variation would be supposed to come from the partial deconfinement of the quarks in a nucleus. As expected from the remarks on the behavior of the potentials as a function of nuclear size for the single Gaussian case, the $T = 0$ phase shift decreases with increasing nucleon size.

Although the ratio of total cross section of nuclei to that of deuterium measured [12, 13] clearly shows a deviation from the predicted ratio with free kaon-nucleon interactions, the energy dependence is not in agreement with the simple change in radius estimated in Ref. [1] being larger at higher incident momenta. Since the model used to calculate the ratio is rather simple, one should consider the possibility that a more realistic model could give the

energy dependence necessary. For this reason we compare the change in cross section that results from a given change in nucleon radius for two different energies. We also make the same type of comparison for changes in interaction strength since that possibility has been suggested as an alternative to an increase in nucleon size[14].

In a series of papers, Caillon and Labarsouque[15] studied this problem with different possibilities for the interaction of the kaon with the nucleus. They are able to partially resolve this problem but have difficulty predicting the correct dependence on atomic number.

In Tables II, III, IV and V is shown the effect on the elastic and total cross sections of the variation in radius and strength. For each isospin the amplitude

$$f_T = \frac{e^{2i\delta_T} - 1}{2i} \quad (37)$$

is calculated and the isospin weighting is done. Since at finite energies a linear expansion of the exponential does not provide a good approximation and since the variations in the two isospin phase shifts with radius and strength go in opposite directions there is not a simple linear proportionality.

From Table II we see that at a beam momentum of about 0.6 GeV/c a change in the radius of 15% corresponds to a change in the differential cross section of 14% and a change in the total cross section of 18%. A change in the radius of 25% gives corresponding changes of 25% and 32%. From Table III we see that at a higher beam momentum the same fractional changes in radii lead to smaller fractional changes in the cross sections. The trend is opposite to what would be needed to give a better fit to the data[12, 13].

Tables IV and V show a similar effect for the variation of the cross sections as a function of the change in the multiplying factor. It is seen that a significant change (in percentage) of the multiplying factor is needed to produce a modest change in the total cross section ratio.

VII. OFF-SHELL AMPLITUDE

From the full solution for the wave function one can calculate the scattering amplitude for off-shell momenta. We have done so at (essentially) zero energy. Since the value of the off-shell amplitude at zero momenta will be equal to the scattering length, dividing this

R(fm)	R/R ⁰	δ_0	δ_1	f_0	f_1	$\frac{3}{4}f_1 + \frac{1}{4}f_0$	$k^2\sigma$	σ/σ^0	σ_T/σ_T^0
0.566	1.00	-0.210	-0.562	$-0.204 + i0.043$	$-0.451 + i0.284$	$-0.389 + i0.224$	0.202	1.00	1.00
0.652	1.15	-0.192	-0.623	$-0.188 + i0.037$	$-0.474 + i0.340$	$-0.402 + i0.264$	0.232	1.15	1.18
0.709	1.25	-0.178	-0.669	$-0.175 + i0.032$	$-0.486 + i0.384$	$-0.409 + i0.296$	0.255	1.26	1.32
0.744	1.31	-0.166	-0.700	$-0.166 + i0.029$	$-0.493 + i0.415$	$-0.411 + i0.318$	0.270	1.34	1.41

TABLE II: Dependence of the s-wave amplitude on the quark radius at $P_{c.m.} = 0.35$ GeV/c ($P_{Lab}=0.599$ GeV/c). R is the rms value of the inter-quark distance in the nucleon. The quantity σ is the (isoscalar) differential cross section which would come from the s-wave alone and the quantity σ_T is the (isoscalar) total cross section from the s-wave alone. The superscript zero refers to the first line in the table i.e. the result obtained from the fit to the data in free space.

R(fm)	R/R ⁰	δ_0	δ_1	f_0	f_1	$\frac{3}{4}f_1 + \frac{1}{4}f_0$	$k^2\sigma$	σ/σ^0	σ_T/σ_T^0
0.566	1.00	-0.312	-0.816	$-0.292 + i0.094$	$-0.499 + i0.531$	$-0.447 + i0.422$	0.378	1.00	1.00
0.652	1.15	-0.287	-0.900	$-0.271 + i0.080$	$-0.487 + i0.614$	$-0.433 + i0.480$	0.418	1.11	1.14
0.709	1.25	-0.267	-0.963	$-0.254 + i0.069$	$-0.469 + i0.673$	$-0.415 + i0.522$	0.445	1.18	1.24
0.744	1.31	-0.253	-1.004	$-0.242 + i0.063$	$-0.453 + i0.711$	$-0.400 + i0.549$	0.462	1.22	1.30

TABLE III: Dependence of the s-wave amplitude on the radius at $P_{c.m.} = 0.5$ GeV/c ($P_{Lab}=0.941$ GeV/c). R is the rms value of the inter-quark distance in the nucleon. The superscript zero refers to the first line in the table i.e. the result obtained from the fit to the data in free space.

amplitude by the scattering length one obtains a function which is unity at $q = 0$. Because the wave function is constant at zero energy the result is equivalent to the Fourier transform of the sum of the four potentials.

Figure 8 shows the result of such a calculation. We see that the half value of the $T=1$ function occurs at around 1 GeV/c. $T=1$ dominates nuclear scattering because of the average weight in the isoscalar average and the relative smallness of the $T=0$ scattering length so its range is perhaps the appropriate one to use.

F	F/F^0	δ_0	δ_1	f_0	f_1	$\frac{3}{4}f_1 + \frac{1}{4}f_0$	σ	σ/σ^0	σ_T/σ_T^0
4.1	1.00	-0.210	-0.562	$-0.204 + i0.043$	$-0.451 + i0.284$	$-0.389 + i0.224$	0.202	1.00	1.00
5.0	1.22	-0.230	-0.591	$-0.222 + i0.052$	$-0.463 + i0.311$	$-0.403 + i0.246$	0.223	1.10	1.10
6.0	1.46	-0.248	-0.618	$-0.238 + i0.060$	$-0.472 + i0.336$	$-0.414 + i0.267$	0.242	1.20	1.19
7.0	1.71	-0.263	-0.640	$-0.251 + i0.068$	$-0.479 + i0.357$	$-0.422 + i0.284$	0.259	1.28	1.27
8.0	1.95	-0.276	-0.659	$-0.262 + i0.074$	$-0.484 + i0.375$	$-0.429 + i0.300$	0.273	1.35	1.34

TABLE IV: Dependence of the s-wave amplitude on the multiplying factor at $P_{c.m.} = 0.35$ GeV/c ($P_{Lab}=0.599$ GeV/c). The superscript zero refers to the first line in the table i.e. the result obtained from the fit to the data in free space.

F	F/F^0	δ_0	δ_1	f_0	f_1	$\frac{3}{4}f_1 + \frac{1}{4}f_0$	$k^2\sigma$	σ/σ^0	σ_T/σ_T^0
4.1	1.00	-0.312	-0.816	$-0.292 + i0.094$	$-0.499 + i0.531$	$-0.447 + i0.422$	0.378	1.00	1.00
5.0	1.22	-0.341	-0.858	$-0.315 + i0.112$	$-0.495 + i0.572$	$-0.450 + i0.457$	0.411	1.08	1.08
6.0	1.46	-0.367	-0.896	$-0.335 + i0.129$	$-0.488 + i0.610$	$-0.450 + i0.489$	0.442	1.17	1.16
7.0	1.71	-0.388	-0.927	$-0.350 + i0.143$	$-0.480 + i0.640$	$-0.447 + i0.516$	0.466	1.23	1.22
8.0	1.95	-0.406	-0.955	$-0.363 + i0.156$	$-0.472 + i0.666$	$-0.444 + i0.538$	0.487	1.29	1.27

TABLE V: Dependence of the s-wave amplitude on the multiplying factor at $P_{c.m.} = 0.50$ GeV/c ($P_{Lab}=0.941$ GeV/c). The superscript zero refers to the first line in the table i.e. the result obtained from the fit to the data in free space.

VIII. DISCUSSION AND CONCLUSIONS

We have calculated the kaon-nucleon scattering amplitude as a function of the nucleon size and interaction strength in the one-gluon-exchange approximation. Two interesting features are observed:

1) The isospin one and isospin zero amplitudes do not behave in the same manner under a change in nucleon size. The isospin zero amplitude actually decreases with increasing nucleon size. Although the potential used is not unique, the fact that the behavior is due




FIG. 8: Ratio of the off-shell to on-shell amplitude at zero energy for $T=1$ (solid) and $T=0$ (dashed). The function falls to 1/2 for $T=1$ at about 1 GeV/c.

to the sharp decrease of η_2 and η_4 with increasing nucleon size leads us to believe that the behavior is more general.

2) A significant renormalization of the potential is required to fit the experimental amplitudes. The factor of 4.1 on the strength (relative to a fixed value of $\gamma = \alpha_s/m_q^2$) of the interaction necessary to bring the quark-gluon exchange into agreement with the s-wave phase shifts is considerable. We have found that the factor needed depends on the extent of the quark distribution of the nucleon and kaon. If this result is confirmed then we must expect additional quark diagrams to play an important role or, alternatively, the exchange of color singlets (for example two-pion exchange with excitation of the K^*) may be important. The very small range of the kaon distribution is also disturbing. Because the potential which results from the fit is very near to that of a hard-sphere, one might think that Pauli blocking might play an important role[17].

We thank Ted Barnes and Eric Swanson for valuable correspondence concerning this work.

This work was supported by the National Science Foundation under contract PHY-0099729.

APPENDIX A: JACOBY COORDINATES

The integrals needed for the calculation are more easily done if the momentum variables are expressed in terms of Jacoby coordinates. Barnes and Swanson[2] use a nucleon wave function represented as

$$\phi(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3) = \frac{3^{\frac{3}{4}}}{\pi^{\frac{3}{2}}\alpha^3} e^{-\frac{\mathbf{p}_1^2 + \mathbf{p}_2^2 + \mathbf{p}_3^2 - \mathbf{p}_1 \cdot \mathbf{p}_2 - \mathbf{p}_2 \cdot \mathbf{p}_3 - \mathbf{p}_3 \cdot \mathbf{p}_1}{3\alpha^2}}. \quad (\text{A1})$$

Expressed in terms of coordinates

$$\mathbf{p} \equiv \mathbf{p}_1 - \mathbf{p}_2; \quad \mathbf{Q} \equiv \frac{\mathbf{p}_1 + \mathbf{p}_2}{2}; \quad \mathbf{q} \equiv \mathbf{p}_3 - \mathbf{Q} \quad (\text{A2})$$

the wave function becomes

$$\phi(\mathbf{p}, \mathbf{q}) = \frac{3^{\frac{3}{4}}}{\pi^{\frac{3}{2}}\alpha^3} e^{-\frac{\mathbf{p}^2}{4\alpha^2}} e^{-\frac{\mathbf{q}^2}{3\alpha^2}}. \quad (\text{A3})$$

We can transform these wave functions into coordinate space. In the Fourier transform we have

$$\mathbf{p}_1 \cdot \mathbf{r}_1 + \mathbf{p}_2 \cdot \mathbf{r}_2 + \mathbf{p}_3 \cdot \mathbf{r}_3 = \mathbf{Q} \cdot \mathbf{R} + \frac{1}{2}\mathbf{p} \cdot \mathbf{r} + \mathbf{q} \cdot \mathbf{r}_3 \quad (\text{A4})$$

where

$$\mathbf{R} = \mathbf{r}_1 + \mathbf{r}_2 + \mathbf{r}_3; \quad \mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2. \quad (\text{A5})$$

Thus the dependence in r-space is

$$\phi(r, r_3) \propto e^{-\frac{\alpha^2 r^2}{4}} e^{-\frac{3\alpha^2 r_3^2}{4}} \quad (\text{A6})$$

giving

$$\langle r^2 \rangle \equiv R^2(N) = 3/\alpha^2 \quad (\text{A7})$$

for the expectation value of the square of the inter-quark separation in the density.

The normalization in this form follows from the normalization given[2]

$$\int d\mathbf{p}_1 d\mathbf{p}_2 d\mathbf{p}_3 \phi^2(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3) \delta(\mathbf{P}_{c.m.} - \mathbf{p}_1 - \mathbf{p}_2 - \mathbf{p}_3) \delta(\mathbf{P}'_{c.m.} - \mathbf{p}_1 - \mathbf{p}_2 - \mathbf{p}_3) \quad (\text{A8})$$

$$= \int d\mathbf{p} d\mathbf{Q} d\mathbf{q} \phi^2(\mathbf{p}, \mathbf{Q}) \delta(\mathbf{P}_{c.m.} - 3\mathbf{Q} - \mathbf{q}) \delta(\mathbf{P}'_{c.m.} - 3\mathbf{Q} - \mathbf{q}) = \frac{\delta(\mathbf{P}_{c.m.} - \mathbf{P}'_{c.m.})}{27} \int d\mathbf{p} d\mathbf{q} \phi^2(p, q).$$

The integral over $\mathbf{p}$ and $\mathbf{q}$ should give 27.

$$\int d\mathbf{p} d\mathbf{q} \phi^2(p, q) = (4\pi)^2 \frac{3^{\frac{3}{2}}}{\pi^3 \alpha^6} \int p^2 dp e^{-\frac{p^2}{2\alpha^2}} \int q^2 dq e^{-\frac{2q^2}{3\alpha^2}} = 27. \quad (\text{A9})$$

The kaon wave function is written in terms of the relative momentum as

$$\phi_{kaon}(p) = \frac{1}{\pi^{\frac{3}{4}} \beta^{\frac{3}{2}}} e^{-\frac{p^2}{8\beta^2}}. \quad (\text{A10})$$

APPENDIX B: DERIVATION OF THE AMPLITUDE EXPRESSIONS

We calculate the expression of Ref. [2] for the case where the initial and final Gaussian parameters are not the same in order to enable a representation of the wave functions as a sum of Gaussian functions.

The strategy is the following:

- a) Rewrite the nucleon wave functions in Jacoby coordinates in the form of Eq. A3.
- b) Transform to r-space to get the proper combinations of the momenta.
- c) Perform the momentum integrals to find 3 δ -functions in $\mathbf{r}$
- d) Carry out the remaining 3 Fourier transforms.

One has two quantities to follow as the integrals are done:

- 1) The imaginary part of the exponent and
- 2) The real parts of the exponent which represent the Gaussian functions and, when the integral is completed, give the dependence of the amplitude in the form of a_i , and b_i .

In this process the factors in front of the exponentials are ignored. They are obtained most easily by setting $\mathbf{k} = \mathbf{k}' = 0$ and performing the momentum space integrals directly. The results for these normalizations (η_i) are given in the summary.

The basic Fourier transform used is given by Eq. 12. A useful transformation is

$$\int d\mathbf{r} e^{i\mathbf{u}\cdot\mathbf{r}} e^{-z_1^2 r^2} e^{-z_2^2 (\mathbf{r}+\mathbf{r}')^2} = e^{-i\xi\mathbf{u}\cdot\mathbf{r}'} e^{-z_{12}^2 r'^2} \int d\mathbf{t} e^{i\mathbf{u}\cdot\mathbf{t}} e^{-2z^2 t^2} \quad (\text{B1})$$

obtained with the change of variables

$$\mathbf{t} = \mathbf{r} + \xi\mathbf{r}', \quad \text{with} \quad \xi = \frac{z_2^2}{z_1^2 + z_2^2} \quad (\text{B2})$$

and

$$z_{12}^2 = \frac{z_1^2 z_2^2}{z_1^2 + z_2^2}; \quad \tilde{z}^2 = \frac{1}{2}(z_1^2 + z_2^2). \quad (\text{B3})$$

If $z_1 = z_2 = z$ then $\xi = \frac{1}{2}$ (this is the case for the same initial and final Gaussian wave functions) the formula simplifies to:

$$e^{-\frac{1}{2}\mathbf{u}\cdot\mathbf{r}'} e^{-\frac{1}{2}z^2 r'^2} \int dt e^{i\mathbf{u}\cdot\mathbf{t}} e^{-2z^2 t^2}. \quad (\text{B4})$$

1. Diagram 1

In this case the integral to be calculated is

$$\begin{aligned} \int d\mathbf{a} d\mathbf{b}_1 d\mathbf{b}_2 \phi_A \left(2\mathbf{a} - \frac{2\rho\mathbf{k}}{1+\rho} \right) \phi_C^* \left(2\mathbf{a} + \frac{2\mathbf{k}'}{1+\rho} - 2\mathbf{k} \right) \phi_B(\mathbf{b}_1, \mathbf{b}_2, -\mathbf{k} - \mathbf{b}_1 - \mathbf{b}_2) \\ \times \phi_D^*(\mathbf{b}_1 + \mathbf{k} - \mathbf{k}', \mathbf{b}_2, -\mathbf{k} - \mathbf{b}_1 - \mathbf{b}_2). \end{aligned} \quad (\text{B5})$$

Here ϕ_A (with Gaussian parameter β_1) and ϕ_C (with Gaussian parameter β_2) represent the two kaon wave functions and ϕ_B (with Gaussian parameter α_1) and ϕ_D (with Gaussian parameter α_2) represent the two nucleon wave functions.

One sees immediately that the kaon wave function integral on $\mathbf{a}$ factors off so that we could do the kaon integral directly in momentum space but to be uniform in procedure we transform to r-space so we have

$$\int d\mathbf{a} e^{-\frac{(\mathbf{a} - \frac{\rho\mathbf{k}}{1+\rho})^2}{2\beta_1^2}} e^{-\frac{(\mathbf{a} + \frac{\mathbf{k}'}{1+\rho} - \mathbf{k})^2}{2\beta_2^2}} \propto \int d\mathbf{a} d\mathbf{r}_5 d\mathbf{r}_6 e^{i(\mathbf{a} - \frac{\rho\mathbf{k}}{1+\rho})\cdot\mathbf{r}_5} e^{-i(\mathbf{a} + \frac{\mathbf{k}'}{1+\rho} - \mathbf{k})\cdot\mathbf{r}_6} e^{-\frac{1}{2}\beta_1^2 r_5^2} e^{-\frac{1}{2}\beta_2^2 r_6^2} \quad (\text{B6})$$

$$\propto e^{i\frac{\mathbf{k}-\mathbf{k}'}{1+\rho}\cdot\mathbf{r}_5} e^{-\tilde{\beta}^2 r_5^2} \propto e^{-\frac{(\mathbf{k}-\mathbf{k}')^2}{4\beta^2(1+\rho)^2}}. \quad (\text{B7})$$

For nucleon B the three momenta $\mathbf{b}_1$, $\mathbf{b}_2$, and $-\mathbf{k} - \mathbf{b}_1 - \mathbf{b}_2$ give for the Jacoby coordinates

$$\mathbf{p} = \mathbf{b}_1 - \mathbf{b}_2, \quad \mathbf{Q} = \frac{\mathbf{b}_1 + \mathbf{b}_2}{2}, \quad \mathbf{q} = -\mathbf{k} - \frac{3}{2}(\mathbf{b}_1 + \mathbf{b}_2) \quad (\text{B8})$$

and hence the B nucleon wave function factor becomes

$$e^{-\frac{(\mathbf{b}_1 - \mathbf{b}_2)^2}{4\alpha_1^2}} e^{-\frac{[\mathbf{k} + \frac{3}{2}(\mathbf{b}_1 + \mathbf{b}_2)]^2}{3\alpha_1^2}}. \quad (\text{B9})$$

For the D nucleon the coordinates $\mathbf{b}_1 + \mathbf{k} - \mathbf{k}'$, $\mathbf{b}_2$, and $-\mathbf{k} - \mathbf{b}_1 - \mathbf{b}_2$ give for Jacoby coordinates

$$\mathbf{p} = \mathbf{k} - \mathbf{k}' + \mathbf{b}_1 - \mathbf{b}_2, \mathbf{Q} = \frac{\mathbf{k} - \mathbf{k}' + \mathbf{b}_1 + \mathbf{b}_2}{2}, \text{ and } \mathbf{q} = -\frac{3}{2}\mathbf{k} + \frac{1}{2}\mathbf{k}' - \frac{3}{2}(\mathbf{b}_1 + \mathbf{b}_2) \quad (\text{B10})$$

and the wave function becomes

$$e^{-\frac{(\mathbf{k}-\mathbf{k}'+\mathbf{b}_1-\mathbf{b}_2)^2}{4\alpha_2^2}} e^{-\frac{[\frac{3}{2}\mathbf{k}-\frac{1}{2}\mathbf{k}'+\frac{3}{2}(\mathbf{b}_1+\mathbf{b}_2)]^2}{3\alpha_2^2}}. \quad (\text{B11})$$

Writing these factors as r-space transforms we have

$$\begin{aligned} & \int d\mathbf{b}_1 d\mathbf{b}_2 d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 e^{i(\mathbf{b}_1-\mathbf{b}_2)\cdot\mathbf{r}_1} e^{-i(\mathbf{k}-\mathbf{k}'+\mathbf{b}_1-\mathbf{b}_2)\cdot\mathbf{r}_2} e^{-\tilde{\alpha}^2(r_1^2+r_2^2)} \\ & \times e^{i[\mathbf{k}+\frac{3}{2}(\mathbf{b}_1+\mathbf{b}_2)]\cdot\mathbf{r}_3} e^{-i[\frac{3}{2}\mathbf{k}-\frac{1}{2}\mathbf{k}'+\frac{3}{2}(\mathbf{b}_1+\mathbf{b}_2)]\cdot\mathbf{r}_4} e^{-\frac{3}{4}\tilde{\alpha}^2(r_3^2+r_4^2)}. \end{aligned} \quad (\text{B12})$$

The integrals on $\mathbf{b}_1$ and $\mathbf{b}_2$ give the delta functions

$$\delta(\mathbf{r}_1 - \mathbf{r}_2 + \frac{3}{2}\mathbf{r}_3 - \frac{3}{2}\mathbf{r}_4), \delta(-\mathbf{r}_1 + \mathbf{r}_2 + \frac{3}{2}\mathbf{r}_3 - \frac{3}{2}\mathbf{r}_4). \quad (\text{B13})$$

We use them to eliminate $\mathbf{r}_2$ and $\mathbf{r}_4$ to find $\mathbf{r}_2 = \mathbf{r}_1$ and $\mathbf{r}_4 = \mathbf{r}_3$.

The integral becomes

$$\int d\mathbf{r}_1 e^{-(\mathbf{k}-\mathbf{k}')\cdot\mathbf{r}_1} e^{-2\tilde{\alpha}^2 r_1^2} \int d\mathbf{r}_3 e^{-i\frac{1}{2}(\mathbf{k}-\mathbf{k}')\cdot\mathbf{r}_3} e^{-\frac{3}{2}\tilde{\alpha}^2 r_3^2} \propto e^{-\frac{(\mathbf{k}-\mathbf{k}')^2}{8\tilde{\alpha}^2}} e^{-\frac{(\mathbf{k}-\mathbf{k}')^2}{24\tilde{\alpha}^2}} \propto e^{-\frac{(\mathbf{k}-\mathbf{k}')^2}{6\tilde{\alpha}^2}}. \quad (\text{B14})$$

Combining with the result for the kaon we have an exponent of

$$(\mathbf{k} - \mathbf{k}')^2 \left[\frac{1}{4\tilde{\beta}^2(1+\rho)^2} + \frac{1}{6\tilde{\alpha}^2} \right] = k^2(1-\mu) \left[\frac{1}{2\tilde{\beta}^2(1+\rho)^2} + \frac{1}{3\tilde{\alpha}^2} \right] \quad (\text{B15})$$

which agrees (in the limit of equal α 's and β 's) with Eq. 41 of Ref. [2]. A numerical check was performed by doing the integral on the momentum $\mathbf{a}$ and then doing the remaining six-dimensional integral by quadrature.

2. Diagram 2

The integral corresponding to diagram 2 is

$$\int d\mathbf{b}_1 d\mathbf{c} d\mathbf{d}_1 \phi_A \left(2\mathbf{c} + \frac{2\mathbf{k}}{1+\rho} - 2\mathbf{k}' \right) \phi_C^* \left(2\mathbf{c} + \frac{2\rho\mathbf{k}'}{1+\rho} \right) \phi_B(\mathbf{b}_1, \mathbf{c}, -\mathbf{k} - \mathbf{b}_1 - \mathbf{c})$$

$$\times \phi_D^*(\mathbf{d}_1, \mathbf{k} - \mathbf{k}' + \mathbf{b}_1 + \mathbf{c} - \mathbf{d}_1, -\mathbf{k} - \mathbf{b}_1 - \mathbf{c}). \quad (\text{B16})$$

Note that there is a typographical error in equation 35 of Ref. [2] where there is a minus sign in front of the term $2\mathbf{k}/(1+\rho)$ in the kaon A wave function [16]. This is only a error of transcription; we agree with all of their final results.

For nucleon B the Jacoby variables are (we leave out $\mathbf{Q}$ which is not needed) $\mathbf{p} = \mathbf{b}_1 - \mathbf{c}$, $\mathbf{q} = -\mathbf{k} - \frac{3}{2}(\mathbf{b}_1 + \mathbf{c})$ and for nucleon D they are $\mathbf{p} = 2\mathbf{d}_1 - \mathbf{k} + \mathbf{k}' - \mathbf{b}_1 - \mathbf{c}$, $\mathbf{q} = -\frac{3}{2}\mathbf{k} + \frac{1}{2}\mathbf{k}' - \frac{3}{2}(\mathbf{b}_1 + \mathbf{c})$. With the vectors $\mathbf{r}_5$ and $\mathbf{r}_6$ being associated with the kaon wave functions the imaginary part of the exponent arising from the Fourier transform is

$$\begin{aligned} & (\mathbf{b}_1 - \mathbf{c}) \cdot \mathbf{r}_1 - [\mathbf{k} + \frac{3}{2}(\mathbf{b}_1 + \mathbf{c})] \cdot \mathbf{r}_2 - [2\mathbf{d}_1 - \mathbf{k} + \mathbf{k}' - \mathbf{b}_1 - \mathbf{c}] \cdot \mathbf{r}_3 + [\frac{3}{2}\mathbf{k} - \frac{1}{2}\mathbf{k}' + \frac{3}{2}(\mathbf{b}_1 + \mathbf{c})] \cdot \mathbf{r}_4. \quad (\text{B17}) \\ & + 2 \left(\mathbf{c} + \frac{\mathbf{k}}{1+\rho} - \mathbf{k}' \right) \cdot \mathbf{r}_5 - 2 \left(\mathbf{c} - \frac{\rho\mathbf{k}'}{1+\rho} \right) \cdot \mathbf{r}_6. \end{aligned}$$

The integrals on $\mathbf{d}_1$, $\mathbf{b}_1$ and $\mathbf{c}$ give $\mathbf{r}_3 = 0$, $\mathbf{r}_2 = \frac{2}{3}\mathbf{r}_1 + \mathbf{r}_4$, and $\mathbf{r}_6 = \mathbf{r}_5 - \mathbf{r}_1$ so that the integral becomes

$$\int d\mathbf{r}_1 d\mathbf{r}_4 d\mathbf{r}_5 e^{i[(-\frac{2}{3}\mathbf{k} - \frac{2\rho\mathbf{k}'}{1+\rho}) \cdot \mathbf{r}_1 + \frac{1}{2}(\mathbf{k} - \mathbf{k}') \cdot \mathbf{r}_4 + 2\frac{(\mathbf{k} - \mathbf{k}')}{1+\rho} \cdot \mathbf{r}_5]} e^{-\alpha_1^2 r_1^2} e^{-\frac{4}{3}\alpha_1^2 (\frac{2}{3}\mathbf{r}_1 + \mathbf{r}_4)^2} e^{-\frac{3}{4}\alpha_2^2 r_4^2} e^{-2\beta_1^2 r_5^2} e^{-2\beta_2^2 (\mathbf{r}_5 - \mathbf{r}_1)^2}. \quad (\text{B18})$$

Defining $\gamma_i^\beta = \beta_i^2/(\beta_1^2 + \beta_2^2)$, we can rewrite the $\mathbf{r}_5$ integral using Eq. B1 as

$$e^{2i\gamma_2^\beta (\frac{\mathbf{k} - \mathbf{k}'}{1+\rho}) \cdot \mathbf{r}_1} e^{-2\beta_{12}^2 r_1^2} \int d\mathbf{s} e^{2i(\frac{\mathbf{k} - \mathbf{k}'}{1+\rho}) \cdot \mathbf{s}} e^{-4\tilde{\beta}^2 s^2}. \quad (\text{B19})$$

Now we do the integral on $\mathbf{r}_4$ using Eq. B1,

$$\int d\mathbf{r}_4 e^{-\frac{1}{2}i(\mathbf{k}' - \mathbf{k}) \cdot \mathbf{r}_4} e^{-\frac{3}{4}\alpha_2^2 r_4^2} e^{-\frac{3}{4}\alpha_1^2 (\frac{2}{3}\mathbf{r}_1 + \mathbf{r}_4)^2} = e^{i\frac{1}{2}(1-\gamma_2^\alpha)(\mathbf{k}' - \mathbf{k}) \cdot \frac{2}{3}\mathbf{r}_1} e^{-\frac{3}{4}\alpha_{12}^2 \frac{4}{9}r_1^2} \int d\mathbf{t} e^{-\frac{1}{2}i(\mathbf{k}' - \mathbf{k}) \cdot \mathbf{t}} e^{-\frac{3}{2}\tilde{\alpha}^2 t^2}. \quad (\text{B20})$$

The contribution of the $\mathbf{s}$ and $\mathbf{t}$ integrals to the exponent is

$$\frac{4(\mathbf{k} - \mathbf{k}')^2}{(1+\rho)^2} \frac{1}{16\tilde{\beta}^2} + \frac{(\mathbf{k} - \mathbf{k}')^2}{4} \frac{2}{12\tilde{\alpha}^2} = \frac{k^2}{2(1+\rho)^2\tilde{\beta}^2} - \frac{\mu k^2}{2(1+\rho)^2\tilde{\beta}^2} + \frac{k^2}{12\tilde{\alpha}^2} - \frac{\mu k^2}{12\tilde{\alpha}^2} \quad (\text{B21})$$

so the contributions of the $\mathbf{s}$ and $\mathbf{t}$ integrals to a_2 and b_2 are

$$\frac{1}{2(1+\rho)^2\tilde{\beta}^2} + \frac{1}{12\tilde{\alpha}^2}; \quad \frac{1}{2(1+\rho)^2\tilde{\beta}^2} + \frac{1}{12\tilde{\alpha}^2}. \quad (\text{B22})$$

The coefficient of $\mathbf{r}_1$ is now

$$\begin{aligned} & -\frac{2}{3}\mathbf{k} - \frac{2\rho\mathbf{k}'}{1+\rho} + 2\gamma_2^\beta \frac{\mathbf{k} - \mathbf{k}'}{1+\rho} + \frac{1}{3}\gamma_1^\alpha(\mathbf{k}' - \mathbf{k}) \\ & = -\frac{[(2 + \gamma_1^\alpha)(1 + \rho) - 6\gamma_2^\beta]\mathbf{k} + [-\gamma_1^\alpha(1 + \rho) + 6(\gamma_2^\beta + \rho)]\mathbf{k}'}{3(1 + \rho)} = C_2\mathbf{k} + C_2'\mathbf{k}' \end{aligned} \quad (\text{B23})$$

where

$$\gamma_i^\alpha = \frac{\alpha_i^2}{\alpha_1^2 + \alpha_2^2}. \quad (\text{B24})$$

The Gaussian in the $\mathbf{r}_1$ integral becomes

$$e^{-(\alpha_1^2 + 2\beta_{12}^2 + \frac{1}{3}\alpha_{12}^2)r_1^2} \quad (\text{B25})$$

so that the contribution of the $\mathbf{r}_1$ integral to the exponential is

$$\frac{(C_2\mathbf{k} + C_2'\mathbf{k}')^2}{4(\alpha_1^2 + 2\beta_{12}^2 + \frac{1}{3}\alpha_{12}^2)} = k^2 \frac{C_2^2 + C_2'^2 + 2C_2C_2'\mu}{4(\alpha_1^2 + 2\beta_{12}^2 + \frac{1}{3}\alpha_{12}^2)}. \quad (\text{B26})$$

so that the contributions to a_2 and b_2 are

$$\frac{C_2^2 + C_2'^2}{4(\alpha_1^2 + 2\beta_{12}^2 + \frac{1}{3}\alpha_{12}^2)}; \quad \text{and} \quad -\frac{C_2C_2'}{2(\alpha_1^2 + 2\beta_{12}^2 + \frac{1}{3}\alpha_{12}^2)}. \quad (\text{B27})$$

A numerical check was done by first calculating the integral on $\mathbf{d}$ in the “p” factor of nucleon D. It can be integrated on freely and gives only a contribution to the normalization of $\pi^{\frac{3}{2}}\alpha_2^3$. The rest of the integral was done by quadrature.

3. Diagram 3

The integral for diagram 3 is

$$\begin{aligned} & \int d\mathbf{a}d\mathbf{b}_2d\mathbf{c}\phi_A\left(2\mathbf{a} - \frac{2\rho\mathbf{k}}{1+\rho}\right)\phi_C^*\left(2\mathbf{c} + \frac{2\rho\mathbf{k}'}{1+\rho}\right)\phi_B(\mathbf{a} - \mathbf{k} + \mathbf{k}', \mathbf{b}_2, -\mathbf{a} - \mathbf{b}_2 - \mathbf{k}') \\ & \quad \times \phi_D^*(\mathbf{a}, \mathbf{b}_2, -\mathbf{a} - \mathbf{b}_2 - \mathbf{k}'). \end{aligned} \quad (\text{B28})$$

The Jacoby variables for nucleon B are: $\mathbf{p} = -\mathbf{k} + \mathbf{k}' + \mathbf{a} - \mathbf{b}_2$; $\mathbf{q} = \frac{1}{2}\mathbf{k} - \frac{3}{2}\mathbf{k}' - \frac{3}{2}(\mathbf{a} + \mathbf{b}_2)$ and for nucleon D: $\mathbf{p} = \mathbf{a} - \mathbf{b}_2$; $\mathbf{q} = -\mathbf{k}' - \frac{3}{2}(\mathbf{a} + \mathbf{b}_2)$, giving the integrand

$$e^{i\left\{(\mathbf{a}-\mathbf{b}_2-\mathbf{k}+\mathbf{k}')\cdot\mathbf{r}_1+\frac{1}{2}[\mathbf{k}-3\mathbf{k}'-3(\mathbf{a}+\mathbf{b}_2)]\cdot\mathbf{r}_2-(\mathbf{a}-\mathbf{b}_2)\cdot\mathbf{r}_3+[\mathbf{k}'+\frac{3}{2}(\mathbf{a}+\mathbf{b}_2)]\cdot\mathbf{r}_4+\left(2\mathbf{a}-\frac{2\rho\mathbf{k}}{1+\rho}\right)\cdot\mathbf{r}_5-\left(2\mathbf{c}+\frac{2\rho\mathbf{k}'}{1+\rho}\right)\cdot\mathbf{r}_6\right\}} \quad (\text{B29})$$

$$\times e^{-\alpha_1^2 r_1^2} e^{-\frac{3}{4}\alpha_1^2 r_2^2} e^{-\alpha_2^2 r_3^2} e^{-\frac{3}{4}\alpha_2^2 r_4^2} e^{-2\beta_1^2 r_5^2} e^{-2\beta_2^2 r_6^2}.$$

Integration on the momenta gives $\mathbf{r}_6 = 0$, $\mathbf{r}_3 = \mathbf{r}_1 + \mathbf{r}_5$ and $\mathbf{r}_2 = \mathbf{r}_4 + \frac{2}{3}\mathbf{r}_5$.

With $\mathbf{z} \equiv \mathbf{k}' - \mathbf{k}$ we have

$$\int d\mathbf{r}_1 d\mathbf{r}_4 d\mathbf{r}_5 e^{i[\mathbf{z} \cdot \mathbf{r}_1 + \frac{1}{2}(\mathbf{k} - 3\mathbf{k}') \cdot (\mathbf{r}_4 + \frac{2}{3}\mathbf{r}_5) + \mathbf{k}' \cdot \mathbf{r}_4 - \frac{2\rho\mathbf{k}}{1+\rho} \cdot \mathbf{r}_5]} e^{-\alpha_1^2 r_1^2} e^{-\frac{3}{4}\alpha_1^2 (\mathbf{r}_4 + \frac{2}{3}\mathbf{r}_5)^2} e^{-\alpha_2^2 (\mathbf{r}_1 + \mathbf{r}_5)^2} e^{-\frac{3}{4}\alpha_2^2 r_4^2} e^{-2\beta_1^2 r_5^2} \quad (\text{B30})$$

$$= \int d\mathbf{r}_1 d\mathbf{r}_4 d\mathbf{r}_5 e^{i\{\mathbf{z} \cdot \mathbf{r}_1 - \frac{1}{2}\mathbf{z} \cdot \mathbf{r}_4 + [\frac{1}{3}(\mathbf{k} - 3\mathbf{k}') - \frac{2\rho\mathbf{k}}{1+\rho}] \cdot \mathbf{r}_5\}} e^{-\alpha_1^2 r_1^2} e^{-\alpha_2^2 (\mathbf{r}_1 + \mathbf{r}_5)^2} e^{-\frac{3}{4}\alpha_2^2 r_4^2} e^{-\frac{3}{4}\alpha_1^2 (\mathbf{r}_4 + \frac{2}{3}\mathbf{r}_5)^2} e^{-2\beta_1^2 r_5^2} \quad (\text{B31})$$

$$= \int d\mathbf{r}_5 \int d\mathbf{t} e^{i\mathbf{z} \cdot \mathbf{t}} e^{-2\tilde{\alpha}^2 t^2} e^{-i\gamma_2^\alpha \mathbf{z} \cdot \mathbf{r}_5} e^{-\alpha_{12}^2 r_5^2} \int d\mathbf{s} e^{-i\frac{1}{2}\mathbf{z} \cdot \mathbf{s}} e^{-\frac{3}{2}\tilde{\alpha}^2 s^2} e^{\frac{1}{3}\gamma_1^\alpha i\mathbf{z} \cdot \mathbf{r}_5} e^{-\frac{1}{3}\alpha_{12}^2 r_5^2} \times e^{i[\frac{1}{3}(\mathbf{k} - 3\mathbf{k}') - \frac{2\rho\mathbf{k}}{1+\rho}] \cdot \mathbf{r}_5} e^{-2\beta_1^2 r_5^2}. \quad (\text{B32})$$

The $\mathbf{s}$ and $\mathbf{t}$ integrals give a contribution to the exponent of

$$\frac{z^2}{8\tilde{\alpha}^2} + \frac{z^2}{24\tilde{\alpha}^2} = \frac{z^2}{6\tilde{\alpha}^2} = \frac{k^2}{3\tilde{\alpha}^2} - \frac{k^2\mu}{3\tilde{\alpha}^2}. \quad (\text{B33})$$

With

$$C_3 \equiv \frac{2[2\gamma_2^\alpha + (2\gamma_2^\alpha - 3)\rho]}{3(1+\rho)}; \quad C'_3 \equiv -\frac{2}{3}(2\gamma_2^\alpha + 1) \quad (\text{B34})$$

the coefficient of $\mathbf{r}_5$ is

$$-\frac{4\gamma_2^\alpha - 1}{3}\mathbf{z} + \frac{\mathbf{k} - 3\mathbf{k}'}{3} - \frac{2\rho\mathbf{k}}{1+\rho} = C_3\mathbf{k} + C'_3\mathbf{k}'. \quad (\text{B35})$$

The $\mathbf{r}_5$ integral is now

$$\int d\mathbf{r}_5 e^{i(C_3\mathbf{k} + C'_3\mathbf{k}') \cdot \mathbf{r}_5} e^{-(\frac{4}{3}\alpha_{12}^2 + 2\beta_1^2)r_5^2} \quad (\text{B36})$$

so its contribution to the exponent is

$$\frac{(C_3\mathbf{k} + C'_3\mathbf{k}')^2}{4(\frac{4}{3}\alpha_{12}^2 + 2\beta_1^2)} = \frac{3k^2(C_3^2 + C_3'^2 + 2C_3C_3'\mu)}{8(2\alpha_{12}^2 + 3\beta_1^2)} \quad (\text{B37})$$

and the contributions to a_3 and b_3 are

$$\frac{3(C_3^2 + C_3'^2)}{8(2\alpha_{12}^2 + 3\beta_1^2)} \quad \text{and} \quad -\frac{3C_3C_3'}{4(2\alpha_{12}^2 + 3\beta_1^2)}. \quad (\text{B38})$$

A numerical check was done by first doing the integral on $\mathbf{c}$ which only appears in kaon C. It can be integrated on freely and gives a contribution to the normalization of $(2\pi)^{\frac{3}{2}}\beta_2^3$.

The remainder of the integral was done by quadrature.

4. Diagram 4

The integral for diagram 4 is

$$\int d\mathbf{a} d\mathbf{b}_1 d\mathbf{c} \phi_A \left(2\mathbf{a} - \frac{2\rho\mathbf{k}}{1+\rho} \right) \phi_C^* \left(2\mathbf{c} + \frac{2\rho\mathbf{k}'}{1+\rho} \right) \phi_B(\mathbf{b}_1, \mathbf{c}, -\mathbf{k} - \mathbf{b}_1 - \mathbf{c}) \\ \times \phi_D^*(\mathbf{k} - \mathbf{k}' - \mathbf{a} + \mathbf{b}_1 + \mathbf{c}, \mathbf{a}, \mathbf{k} - \mathbf{b}_1 - \mathbf{c}). \quad (\text{B39})$$

For the Jacoby variables for nucleon B we have: $\mathbf{p} = \mathbf{b}_1 - \mathbf{c}$; $\mathbf{q} = -\mathbf{k} - \frac{3}{2}(\mathbf{b}_1 + \mathbf{c})$ and for nucleon D: $\mathbf{p} = \mathbf{k} - \mathbf{k}' - 2\mathbf{a} + \mathbf{b}_1 + \mathbf{c}$; $\mathbf{q} = -\frac{3}{2}\mathbf{k} + \frac{1}{2}\mathbf{k}' - \frac{3}{2}(\mathbf{b}_1 + \mathbf{c})$.

The imaginary part of the exponent is

$$i \left\{ (\mathbf{b}_1 - \mathbf{c}) \cdot \mathbf{r}_1 - \left[\mathbf{k} + \frac{3}{2}(\mathbf{b}_1 + \mathbf{c}) \right] \cdot \mathbf{r}_2 - (\mathbf{k} - \mathbf{k}' - 2\mathbf{a} + \mathbf{b}_1 + \mathbf{c}) \cdot \mathbf{r}_3 \right. \\ \left. - \left[-\frac{3}{2}\mathbf{k} + \frac{1}{2}\mathbf{k}' - \frac{3}{2}(\mathbf{b}_1 + \mathbf{c}) \right] \cdot \mathbf{r}_4 + \left(2\mathbf{a} - \frac{2\rho\mathbf{k}}{1+\rho} \right) \cdot \mathbf{r}_5 - \left(2\mathbf{c} - \frac{2\rho\mathbf{k}'}{1+\rho} \right) \cdot \mathbf{r}_6 \right\}. \quad (\text{B40})$$

The $\mathbf{a}$, $\mathbf{b}_1$ and $\mathbf{c}$ integrals give $\mathbf{r}_3 = -\mathbf{r}_5$, $\mathbf{r}_6 = -\mathbf{r}_1$ and $\mathbf{r}_2 = \frac{2}{3}\mathbf{r}_1 + \frac{2}{3}\mathbf{r}_5 + \mathbf{r}_4$.

The Gaussian factors are

$$e^{-(\alpha_1^2+2\beta_2^2)r_1^2} e^{-(\alpha_2^2+2\beta_1^2)r_5^2} e^{-\frac{3}{4}\alpha_2^2 r_4^2} e^{-\frac{3}{4}\alpha_1^2 (\frac{2}{3}\mathbf{r}_1 + \frac{2}{3}\mathbf{r}_5 + \mathbf{r}_4)^2} \quad (\text{B41})$$

and the imaginary part of the exponent

$$-2 \left(\frac{1}{3}\mathbf{k} + \frac{\rho\mathbf{k}'}{1+\rho} \right) \cdot \mathbf{r}_1 + \frac{1}{2}(\mathbf{k} - \mathbf{k}') \cdot \mathbf{r}_4 + \left(\frac{1}{3}\mathbf{k} - \mathbf{k}' - \frac{2\rho\mathbf{k}}{1+\rho} \right) \cdot \mathbf{r}_5 \equiv \mathbf{u}_1 \cdot \mathbf{r}_1 + \mathbf{u}_4 \cdot \mathbf{r}_4 + \mathbf{u}_5 \cdot \mathbf{r}_5. \quad (\text{B42})$$

We make the transformation

$$\mathbf{s} = \mathbf{r}_1 + \mathbf{r}_5; \quad \mathbf{t} = \mathbf{r}_1 - \frac{w^2}{v^2}\mathbf{r}_5; \quad \mathbf{r}_1 = (v^2\mathbf{t} + w^2\mathbf{s})/(v^2 + w^2); \quad \mathbf{r}_5 = v^2(\mathbf{s} - \mathbf{t})/(v^2 + w^2) \quad (\text{B43})$$

where

$$v^2 = \alpha_1^2 + 2\beta_2^2; \quad w^2 = \alpha_2^2 + 2\beta_1^2. \quad (\text{B44})$$

Now

$$v^2 r_1^2 + w^2 r_5^2 = \frac{v^2}{v^2 + w^2} (v^2 t^2 + w^2 s^2) \quad (\text{B45})$$

and the integral becomes

$$\int d\mathbf{s} d\mathbf{t} d\mathbf{r}_4 e^{i \left[\frac{v^2(\mathbf{u}_1 - \mathbf{u}_5) \cdot \mathbf{t}}{v^2 + w^2} + \frac{(w^2\mathbf{u}_1 + v^2\mathbf{u}_5) \cdot \mathbf{s}}{v^2 + w^2} + \mathbf{u}_4 \cdot \mathbf{r}_4 \right]} e^{-(v^2 + w^2)t^2} e^{-\frac{v^2 w^2 s^2}{v^2 + w^2}} e^{-\frac{3}{4}\alpha_2^2 r_4^2} e^{-\frac{3}{4}\alpha_1^2 (\frac{2}{3}\mathbf{s} + \mathbf{r}_4)^2}. \quad (\text{B46})$$

The integral on $\mathbf{t}$ can now be done to give a contribution to the exponent of

$$\frac{v^4(\mathbf{u}_1 - \mathbf{u}_5)^2}{(v^2 + w^2)^2} \frac{(v^2 + w^2)}{4v^4} = \frac{(\mathbf{k}' - \mathbf{k})^2(1 - \rho)^2}{8(\tilde{\alpha} + 2\tilde{\beta})(1 + \rho)^2}. \quad (\text{B47})$$

The $\mathbf{r}_4$ integral becomes

$$\int d\mathbf{r}_4 e^{i\mathbf{u}_4 \cdot \mathbf{r}_4} e^{-\frac{3}{4}\alpha_2^2 r_4^2} e^{-\frac{3}{4}\alpha_1^2 (\frac{2}{3}\mathbf{s} + \mathbf{r}_4)^2} = e^{-i\frac{2}{3}\gamma_1^\alpha \mathbf{u}_4 \cdot \mathbf{s}} e^{-\frac{3}{4}\alpha_{12}^2 (\frac{2}{3})^2 s^2} \int d\mathbf{v} e^{i\mathbf{u}_4 \cdot \mathbf{v}} e^{-\frac{3}{2}\tilde{\alpha}^2 v^2} \quad (\text{B48})$$

and the $\mathbf{v}$ integral gives

$$\frac{(\mathbf{k}' - \mathbf{k})^2}{4} \frac{2}{12\tilde{\alpha}^2} = \frac{(\mathbf{k}' - \mathbf{k})^2}{24\tilde{\alpha}^2}. \quad (\text{B49})$$

The $\mathbf{s}$ integral becomes

$$\int d\mathbf{s} e^{i \left[\frac{w^2 \mathbf{u}_1 + v^2 \mathbf{u}_5}{v^2 + w^2} - \frac{2}{3} \gamma_1^\alpha \mathbf{u}_4 \right] \cdot \mathbf{s}} e^{-\left(\frac{v^2 w^2}{v^2 + w^2} + \frac{1}{3} \alpha_{12}^2 \right) s^2} \quad (\text{B50})$$

$$\frac{w^2 \mathbf{u}_1 + v^2 \mathbf{u}_5}{v^2 + w^2} - \frac{2}{3} \gamma_1^\alpha \mathbf{u}_4 = - \frac{\left[2w^2 - v^2 + \frac{6v^2 \rho}{1 + \rho} + (v^2 + w^2) \gamma_1^\alpha \right] \mathbf{k} + \left[\frac{6b^2 \rho}{1 + \rho} + 3v^2 - (v^2 + w^2) \gamma_1^\alpha \right] \mathbf{k}'}{3(v^2 + w^2)} \quad (\text{B51})$$

$$= C_4 \mathbf{k} + C'_4 \mathbf{k}'.$$

so the contribution to the exponent from the $\mathbf{s}$ integral is

$$\frac{(C_4 \mathbf{k} + C'_4 \mathbf{k}')^2}{4 \left(\frac{v^2 w^2}{v^2 + w^2} + \frac{1}{3} \alpha_{12}^2 \right)} = \frac{k^2 (C_4^2 + C_4'^2 + 2C_4 C'_4 \mu)}{4 \left(\frac{v^2 w^2}{v^2 + w^2} + \frac{1}{3} \alpha_{12}^2 \right)} \quad (\text{B52})$$

and the contribution to a_4 and b_4 are

$$\frac{C_4^2 + C_4'^2}{4 \left(\frac{v^2 w^2}{v^2 + w^2} + \frac{1}{3} \alpha_{12}^2 \right)} \quad \text{and} \quad - \frac{C_4 C'_4}{2 \left(\frac{v^2 w^2}{v^2 + w^2} + \frac{1}{3} \alpha_{12}^2 \right)}. \quad (\text{B53})$$

For a numerical check, the $\mathbf{a}$ integral which only affects kaon A and the “ $\mathbf{p}$ ” factor of nucleon D was done. With $\mathbf{x} = \mathbf{a} - \frac{\rho \mathbf{k}}{1 + \rho}$ we have

$$\begin{aligned} \int d\mathbf{a} e^{-\frac{(\mathbf{a} - \frac{\rho \mathbf{k}}{1 + \rho})^2}{2\beta_1^2}} e^{-\frac{(\mathbf{k} - \mathbf{k}' - 2\mathbf{a} + \mathbf{b} + \mathbf{c})^2}{4\alpha_2^2}} &= \int d\mathbf{x} e^{-\frac{x^2}{2\beta_1^2}} e^{-\frac{[\mathbf{x} + \frac{\rho \mathbf{k}}{1 + \rho} - \frac{1}{2}(\mathbf{k} - \mathbf{k}' + \mathbf{b} + \mathbf{c})]^2}{\alpha_2^2}} \\ &= \frac{\sqrt{\pi}}{4} \left(\frac{2\alpha_2^2 \beta_1^2}{\alpha_2^2 + 2\beta_1^2} \right)^{\frac{3}{2}} e^{\frac{[\frac{\rho \mathbf{k}}{1 + \rho} - \frac{1}{2}(\mathbf{k} - \mathbf{k}' + \mathbf{b} + \mathbf{c})]^2}{\alpha_2^2 + 2\beta_1^2}}. \end{aligned} \quad (\text{B54})$$

With this first integral done, the remaining 6-dimensional integral was done numerically.

[1] P. B. Siegel, W. B. Kaufmann and W. R. Gibbs, Phys. Rev. C **31**, 2184 (1985)

- [2] T. Barnes and E. S. Swanson, Phys. Rev. C **49**, 1166 (1994)
- [3] T. Barnes, N. Black and E. S. Swanson, Phys. Rev. C **63**, 025204 (2001)
- [4] T. E. O. Ericson and M. Rosa-Clot, Nucl. Phys. **A405**, 497 (1983).
- [5] J. L. Friar, B. F. Gibson and G. L. Payne, Phys. Rev. C **30**, 1084 (1984).
- [6] J. L. Ballot and M. R. Robilotta, Phys. Rev. C **45**, 986 (1992); Phys. Rev. C **45**, 990 (1992);
J. L. Ballot, A. M. Eiró and M. R. Robilotta, Phys. Rev. C **40**, 1459 (1989).
- [7] W. R. Gibbs and R. Arceo, Phys. Rev. C **75**, 035204 (2007)
- [8] N. Black, J. Phys. G **28**, 1953 (2002)
- [9] M. Lissia, M.-C. Chu, J. W. Negele and J. M. Grandy, Nucl. Phys. **A555**, 272 (1993).
- [10] C. Alexandrou, Ph. de Forcrand and A Tsapalis, Phys. Rev. D **68**, 074504 (2003)
- [11] J. P. Hilbert, N. Black, T. Barnes, and E. S. Swanson, Phys. Rev. C **75**, 064907 (2007)
- [12] R. A. Krauss, J. Alster, D. Ashery, S. Bart, R. E. Chrien, J. C. Hiebert, R. R. Johnson, T. Kishimoto, I. Mardor, Y. Mardor, M. A. Moinester, R. Olshevsky, E. Piasetzky, P. H. Pile, R. Sawafta, R. L. Stearns, R. J. Sutter, R. Weiss, and A. I. Yavin, Phys. Rev. C **46**, 655 (1992)
- [13] R. Weiss, J. Aclander, J. Alster, M. Barakat, S. Bart, R. E. Chrien, R. A. Krauss, K. Johnston, I. Mardor, Y. Mardor, S. May Tal-beck, E. Piasetzky, P. H. Pile, R. Sawafta, H. Seyfarth, R. L. Stearns, R. J. Sutter, and A. I. Yavin, Phys. Rev. C **49**, 2569 (1994)
- [14] G.E. Brown, C.B. Dover, P.B. Siegel and W. Weise, Phys. Rev. Lett. **60**, 2723 (1988)
- [15] J. C. Caillon and J. Labarsouque, Phys. Rev. C **53**, 1993 (1996); Nucl. Phys. **A589**, 609 (1995); Phys. Rev. C **45**, 2503 (1992)
- [16] Private communication, Eric Swanson
- [17] I. Bender, H. G. Dosch, H. J. Pirner, and H. G. Kruse, Nucl. Phys. **A414**, 359 (1984)